

Embryogenesis of an Invertebrate Lymphoid Organ: The *Asterina Gibbosa* Axial Organ (Asterids, Echinodermata)

Michel Leclerc*

Immunology of Invertebrates 556 rue Isabelle Romée 45640 Sandillon (France)

***Corresponding Author:** Michel Leclerc, Immunology of Invertebrates 556 rue Isabelle Romée 45640 Sandillon (France)

Abstract

The metamerization of the coelom leads to the axial complex genesis. Mesodermic cells appear besides the stone canal (larva 12 days old): they constitute the axial organ (AO) which develop, in the ventral side, the oral part (PO). This last part corresponds (sensu stricto) to the ancestral lymphoid organ.

Keywords: Echinodermata, Asterids, axial organ, mesodermic cells

Introduction

In 1896, Mac Bride (Ref.1) described the development of the Asterid : *Asterina gibbosa*, and particularly the development of the axial organ (AO): an ancestral lymphoid organ (Leclerc, Ref.2).

We recall in this work, the different steps of the embryogenesis of *Asterina gibbosa*

Material and methods

Sea star larva were obtained in our laboratory, in aquarum, in running sea water, at 11°C, in June of each year, after spawning . Just before this last one we used the "1 methyl-Adenin" which facilitates it (injection at low concentration: 10µl of 1 methyl-Adenin at a concentration of 1 mg/ml, in the coelomic cavity of the adult *Asterina gibbosa*).

Young sea stars (1 month old) were kept in another aquarum and used for our experiments.

Histologic Studies

Larva and young sea stars, were fixed in Carnoy, dehydrated from alcohol 30° to alcohol 100 °.

Before embedding in Paraffin, animals were placed in a solution of Xylen. For 30 min.

Cuts of 4µ were realized with a classical microtome.

Coloration of slides were performed with either Hemalun or Toluidin-Blue Eosin.


Observations were done with a Light Microscope.

Results

Figure 28 shows the "birth" of the axial organ. Many mesodermic cells surround the stone canal (CS). A primordial gonocyte (Gp) (Ref.3) can be observed next to the mesenchymatous cells. It seems that the aboral part (Pa) appears before the oral part (Po) of the axial organ (Ref 3) during embryogenesis in larva 12 days old. At 15-16 days old the larva becomes a young sea star.

Figure 31 shows clearly the stone canal and the oral part (Po) which corresponds to the main lymphoid organ "sensu stricto". We observe in this young sea star (1 month old), nucleus of cells which'll become the future lymphocytes

Embryogenesis of an Invertebrate Lymphoid Organ: The Asterina Gibbosa Axial Organ (Asterids, Echinodermata)



Discussion and Conclusion

The embryogenesis of the sea star *Asterina gibbosa* can be summarized as following:

- 1) Metamerization of the coelom which leads to the mesodermic formation of the axial organ besides the stone canal, at the beginning of the larva stages (From larva 10 days old to 15 days old).
- 2) Formation of the oral part of the axial organ in larva 15 days old. This part constitutes the future

lymphoid organ with its humoral potentialities (Ref 4) which are found in adult animals.

Reference

- [1] Mac Bride, E.W (1896) Quart.J.Micr.Sci., 38:339-411
- [2] Leclerc, M. Et al (1980) Dev. Comp. Immunol., 4:605-615
- [3] Leclerc, M (1974) Thèse de Doctorat ès Sciences, Orléans (France)
- [4] Vincent, N. Et al ((2013) Meta Gene 2:320-322

Citation: Michel Leclerc. (2020), "Embryogenesis of an Invertebrate Lymphoid Organ: The Asterina Gibbosa Axial Organ (Asterids, Echinodermata)", Arch Health Sci; 4(1): 1-2.

DOI: 10.31829/2641-7456/ahs2020-4(1)-116

Copyright: © 2020 Michel Leclerc. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.